

12th International Vilnius Conference on Probability Theory and Mathematical Statistics and  
 2018 IMS Annual Meeting on Probability and Statistics  
 Abstracts, p. 321  
 © 2018 VTeX, Vilnius

## CALIBRATED NONPARAMETRIC BOOTSTRAP APPROXIMATIONS OF FINITE POPULATION $L$ -STATISTICS

ANDRIUS ČIGINAS

Vilnius University, Vilnius, Lithuania  
 e-mail: [andrius.ciginas@mii.vu.lt](mailto:andrius.ciginas@mii.vu.lt)

DALIUS PUMPUTIS

Vilnius Gediminas Technical University, Vilnius, Lithuania  
 e-mail: [dalius.pumputis@vgtu.lt](mailto:dalius.pumputis@vgtu.lt)

Consider a study variable  $x$  taking real values in  $\mathcal{X} = \{x_1, \dots, x_N\}$  in the population  $\mathcal{U} = \{1, \dots, N\}$ ;  $\mathbb{X} = \{X_1, \dots, X_n\}$  is the set of measurements of the simple random sample units  $\{1, \dots, n\}$ ,  $n < N$ , drawn without replacement from  $\mathcal{U}$ . The  $L$ -statistic

$$L = L_n(\mathbb{X}) = \frac{1}{n} \sum_{j=1}^n c_{j,n} X_{j:n}$$

is a linear combination of the order statistics  $X_{1:n} \leq \dots \leq X_{n:n}$  of  $\mathbb{X}$  with the real coefficients  $c_{j,n} = J(j/(n+1))$ , where  $J: (0, 1) \rightarrow \mathbb{R}$ , called weights. The sample mean, Gini's mean difference statistic and trimmed means are particular cases of  $L$ .

We aim to estimate the distribution function

$$F_S(y) = \mathbb{P}\{\hat{\sigma}_J^{-1}(L - \mathbb{E}L) \leq y\}$$

of the Studentized  $L$ -statistic, where

$$\hat{\sigma}_J^2 = \hat{\sigma}_J^2(\mathbb{X}) = \left(1 - \frac{n}{N}\right) \frac{n-1}{n} \sum_{k=1}^n (L_{(k)} - \bar{L})^2, \quad \bar{L} = \frac{1}{n} \sum_{k=1}^n L_{(k)},$$

is the jackknife estimator of the variance  $\sigma^2 = \text{Var}L$ . Here  $L_{(k)} = L_{n-1}(\mathbb{X} \setminus \{X_k\})$ ,  $1 \leq k \leq n$ , are  $L$ -statistics with the weights  $c_{j,n-1} = J(j/n)$ ,  $1 \leq j \leq n-1$ .

Usually, empirical Edgeworth expansions (EEE) and bootstrap approximations improve the normal approximation applied to  $F_S(y)$ , if the sample size is not large enough. If well-correlated auxiliary information is available for all units of  $\mathcal{U}$ , the calibration technique [1] corrects EEEs based on the sample  $\mathbb{X}$  only [2]. We construct a new calibrated nonparametric bootstrap approximation to  $F_S(y)$ , and compare it with selected calibrated (parametric) EEEs and some other approximations in a simulation study.

### References

1. Deville, J.C. and Särndal, C.-E. (1992). Calibration estimators in survey sampling. *Journal of the American Statistical Association*, 87: 376–382. [MR1173804](#)
2. Pumputis, D. and Čiginas, A. (2013). Estimation of parameters of finite population  $L$ -statistics. *Nonlinear Analysis: Modelling and Control*, 18: 327–343. [MR3072937](#)